

MALNORMAL SUBGROUPS OF LATTICES AND THE PUKÁNSZKY INVARIANT IN GROUP FACTORS

GUYAN ROBERTSON AND TIM STEGER

ABSTRACT. Let G be a connected semisimple real algebraic group. Assume that $G(\mathbb{R})$ has no compact factors and let Γ be a torsion-free uniform lattice subgroup of $G(\mathbb{R})$. Then Γ contains a malnormal abelian subgroup A . This implies that the II_1 factor $\text{VN}(\Gamma)$ contains a masa $\mathfrak{A}$ with Pukánszky invariant $\{\infty\}$.

1. INTRODUCTION

A subgroup Γ_0 of a group Γ is *malnormal* if $x\Gamma_0x^{-1} \cap \Gamma_0 = \{1\}$ for all $x \in \Gamma - \Gamma_0$. An abelian malnormal subgroup is necessarily maximal abelian. The main result of this article is Theorem 1.1, which rests upon work of Prasad and Rapinchuk [PrR].

Theorem 1.1. *Let G be a connected semisimple real algebraic group and let d be the $\mathbb{R}$ -rank of G . Assume that $G(\mathbb{R})$ has no compact factors and let Γ be a torsion-free uniform lattice subgroup of $G(\mathbb{R})$. Then Γ contains a malnormal abelian subgroup $A \cong \mathbb{Z}^d$.*

Theorem 1.1 will be applied to the group factor $\text{VN}(\Gamma)$. Recall that if Γ is a group, then the von Neumann algebra $\text{VN}(\Gamma)$ is the convolution algebra

$$\text{VN}(\Gamma) = \{f \in \ell^2(\Gamma) : f \star \ell^2(\Gamma) \subseteq \ell^2(\Gamma)\}.$$

It is well known that if Γ is an infinite conjugacy class group then $\text{VN}(\Gamma)$ is a factor of type II_1 . This is true if Γ is a lattice in a semisimple Lie group [GHJ, Lemma 3.3.1]. If Γ_0 is a subgroup of Γ , then $\text{VN}(\Gamma_0)$ embeds naturally as a subalgebra of $\text{VN}(\Gamma)$ via $f \mapsto \overline{f}$, where

$$\overline{f}(x) = \begin{cases} f(x) & \text{if } x \in \Gamma_0, \\ 0 & \text{otherwise.} \end{cases}$$

This article is concerned with examples where $\Gamma_0 = A$ is an abelian subgroup of Γ and $\mathfrak{A} = \text{VN}(A)$ is a maximal abelian $\star$ -subalgebra (masa) of $\mathfrak{M} = \text{VN}(\Gamma)$. Recall that $\mathfrak{A}$ is the von Neumann subalgebra of $B(\ell^2(\Gamma))$ defined by the left convolution operators

$$\lambda(f) : \phi \mapsto f \star \phi$$

Date: July 17, 2009.

2000 Mathematics Subject Classification. 22D25, 22E40, 20G20.

where $f \in \ell^2(A)$ and $f \star \ell^2(\Gamma) \subseteq \ell^2(\Gamma)$. The algebra $\mathfrak{A}$ also acts on $\ell^2(\Gamma)$ by right convolution

$$\rho(f) : \phi \mapsto \phi \star f.$$

Let $\mathfrak{A}^{opp}$ be the von Neumann subalgebra of $B(\ell^2(\Gamma))$ defined by this right action of $\mathfrak{A}$. Let $\mathfrak{B}$ be the von Neumann subalgebra of $B(\ell^2(\Gamma))$ generated by $\mathfrak{A} \cup \mathfrak{A}^{opp}$ and let p denote the orthogonal projection of $\ell^2(\Gamma)$ onto the closed subspace generated by $\mathfrak{A}$. Then p is in the centre of the commutant $\mathfrak{B}'$, and $\mathfrak{B}'p$ is abelian. The von Neumann algebra $\mathfrak{B}'(1 - p)$ is of type I and may therefore be expressed as a direct sum $\mathfrak{B}_{n_1} \oplus \mathfrak{B}_{n_2} \oplus \dots$ of algebras $\mathfrak{B}_{n_i}$ of type I_{n_i} , where $1 \leq n_1 < n_2 < \dots \leq \infty$. The *Pukánszky invariant* [SS2, Chapter 7] is the set $\{n_1, n_2, \dots\}$. It is an isomorphism invariant of the pair $(\mathfrak{A}, \mathfrak{M})$. It has been shown [NeS, Corollary 3.3] that each nonempty subset S of the natural numbers containing 1 can be realized as the Pukánszky invariant of some masa in the hyperfinite II_1 factor R . This was extended [SS1, DSS] to subsets S containing ∞ for R and for the free group factor. It was later extended [Whi] to arbitrary subsets $S \subset \mathbb{N} \cup \{\infty\}$ for R (and for certain other McDuff factors).

It is known that every factor of type II_1 contains a singular masa [Po1]. S. Popa [Po2, Remark 3.4] showed that if the Pukánszky invariant of $(\mathfrak{A}, \mathfrak{M})$ does not contain 1, then $\mathfrak{A}$ is a singular masa in $\mathfrak{M}$. K. Dykema [Dyk] has shown (using Voiculescu's free entropy dimension) that the Pukánszky invariant of any masa in the free group factor must either contain ∞ or be unbounded. This means that it is not possible for any singleton other than $\{\infty\}$ to be a possible Pukánszky invariant occurring in every II_1 factor. Jolissaint [Jol] has shown that if F_0 is the cyclic subgroup generated by the first generator of Thompson's group F then $\text{VN}(F_0)$ has Pukansky invariant $\{\infty\}$. A natural question arises.

- Does every II_1 factor $\mathfrak{M}$ contain a masa $\mathfrak{A}$ with Pukánszky invariant $\{\infty\}$?

This article uses Theorem 1.1 to provide an affirmative answer for $\mathfrak{M} = \text{VN}(\Gamma)$, where Γ is a torsion-free uniform lattice subgroup of a connected semisimple real algebraic group G without compact factors. If G has $\mathbb{R}$ -rank ≥ 2 , then $\text{VN}(\Gamma)$ has Kazhdan's property T . This is the first result on possible values of the Pukánszky invariant in a II_1 factor with property T .

Many thanks are due to G. Prasad and Y. Shalom for their assistance with the proofs of Theorem 1.1 and Theorem 3.2 respectively. S. White provided valuable background information.

2. MALNORMAL ABELIAN SUBGROUPS OF LATTICES

This section is devoted to the proof of Theorem 1.1. Let G be a connected semisimple real algebraic group and let d be the $\mathbb{R}$ -rank of G . Assume that $G(\mathbb{R})$ has no compact factors and let Γ be a torsion-free uniform lattice subgroup of $G(\mathbb{R})$.

Since Γ is finitely generated, $\Gamma < G(\mathbb{K})$ for some finitely generated subfield $\mathbb{K}$ of $\mathbb{R}$. By the Borel Density Theorem [Mar, Chapter II, Corollary 4.4], Γ is Zariski dense in $G(\mathbb{R})$. Therefore, according to Theorems 1 and 2 of [PrR], there exists a maximal abelian torus subgroup T of G with the following properties.

- (1) The $\mathbb{R}$ -rank of T is d .
- (2) $A = T(\mathbb{R}) \cap \Gamma$ is a uniform lattice in $T(\mathbb{R})$.
- (3) T has no proper algebraic subgroups defined over $\mathbb{K}$.

Moreover, T is the $\mathbb{K}$ -Zariski closure of a single $\mathbb{R}$ -regular element x_0 in A [PrR]. In fact there are many such elements x_0 [PrR, Remark 2]. We claim that A is a malnormal subgroup of Γ . To this end, fix an arbitrary element $x \in \Gamma - A$. We must show that $xAx^{-1} \cap A = \{1\}$.

Let T_s (respectively T_a) be the maximal $\mathbb{R}$ -split (respectively $\mathbb{R}$ -anisotropic) subtorus of T . Then $T = T_s \cdot T_a$ (an almost direct product) [Bor, Proposition 8.15] and T_s is a maximal $\mathbb{R}$ -split torus in G [PrR, Remark 1]. Thus $T(\mathbb{R}) = S \cdot C$, where $S = T_s(\mathbb{R}) \cong \mathbb{R}^d$ and $C = T_a(\mathbb{R}) \cong (\mathbb{R}/\mathbb{Z})^r$, where r is the dimension of T_a . Since Γ is torsion free and discrete, A is a uniform lattice in S . In particular, $A \cong \mathbb{Z}^d$.

Since T is the $\mathbb{K}$ -Zariski closure of A , it follows that T is defined over $\mathbb{K}$. Since $x \in \Gamma \subseteq G(\mathbb{K})$, xTx^{-1} is also defined over $\mathbb{K}$. According to condition (3), there are only two possibilities:

- $T \cap xTx^{-1} = \{1\}$;
- $T = xTx^{-1}$.

In the first case we also have $T(\mathbb{K}) \cap xT(\mathbb{K})x^{-1} = \{1\}$, and *a fortiori* $A \cap xAx^{-1} = \{1\}$, as required.

To show that the second case does not occur, assume that $T = xTx^{-1}$. This implies that $T(\mathbb{R})$ is stable under conjugation by x . Also Γ is stable under conjugation by x . Therefore $xAx^{-1} = A$, since $A = \Gamma \cap T(\mathbb{R})$. There are two possibilities to consider for the action $\alpha_x : a \mapsto xax^{-1}$ on A .

- (a) α_x fixes only the trivial element of A ;
- (b) α_x fixes some nontrivial element of A .

Since conjugation by x stabilizes T , it also stabilizes T_s and T_a separately [Bor, Proposition 8.15(3)]. Thus $xsx^{-1} = S$. The symmetric space of $G(\mathbb{R})$ is $X = G(\mathbb{R})/K$, where K is a maximal compact subgroup of $G(\mathbb{R})$. The group Γ acts freely on X , since it is torsion free. Since T_s is a maximal $\mathbb{R}$ -split torus of G , there is a unique flat F in X

such that $SF = F$ and S acts simply transitively on F [Mos, Lemma 5.1]. Now xF is another such flat, since

$$SxF = (xSx^{-1})xF = xSF = xF.$$

Hence $xF = F$.

The action of x on F is by some rigid motion and the action of A on F is by translations. No nontrivial element in Γ can act trivially on F , so we can calculate the conjugation by x of any element $y \in A$ by considering the actions of these elements on F . The two cases above correspond to:

- (a) x acts on F by a rigid motion whose linear part has trivial 1-eigenspace;
- (b) x acts on F by a rigid motion whose linear part has nontrivial 1-eigenspace.

In case (a), x necessarily has a fixed point in F . Therefore $x = 1$, since Γ acts freely on X .

Consider case (b). The algebraic subgroup $Z_G(x) \cap T$, which consists of the elements commuting with x , is defined over $\mathbb{K}$. In case (b), $Z_G(x) \cap T$ contains nontrivial elements of A . That is, it has nontrivial $\mathbb{K}$ -points. Hence, it must be nontrivial (as an algebraic group). By condition (3) it must be all of T . Hence x commutes with every element of T . Therefore the algebraic closure of $\{x\} \cup T$ over $\mathbb{K}$ is commutative. However T is a maximal abelian subgroup over $\mathbb{K}$, and so $x \in T(\mathbb{K})$. Therefore $x \in A$, contrary to assumption. This completes the proof of Theorem 1.1.

3. THE PUKÁNSZKY INVARIANT

The following result was proved in [RoS, Proposition 3.6] and later extended in [SS1, Theorem 4.1].

Proposition 3.1. *Suppose that A is an abelian subgroup of a countable group Γ such that $\mathfrak{A} = \text{VN}(A)$ is a masa of $\text{VN}(\Gamma)$. If A is malnormal in Γ then the Pukánszky invariant of $\mathfrak{A}$ contains precisely one element $n = \#(A \backslash \Gamma / A - \{A\})$.*

In view of Theorem 1.1, the next result is enough to provide examples of masas with Pukánszky invariant $\{\infty\}$.

Theorem 3.2. *Let G be a connected semisimple real algebraic group. Assume that $G(\mathbb{R})$ has no compact factors. Let Γ be a torsion free uniform lattice subgroup of $G(\mathbb{R})$ and let $A < \Gamma$ be an abelian subgroup. Then*

$$\#(A \backslash \Gamma / A) = \infty.$$

Proof. Suppose that $\#(A \backslash \Gamma / A) < \infty$. Then

$$\Gamma = \bigcup_{x \in F} AxA$$

where $F \subset \Gamma$ is finite. Taking Zariski closures, it follows from the Borel Density Theorem that

$$(1) \quad G(\mathbb{R}) = \bar{\Gamma} = \bigcup_{x \in F} \overline{Ax\bar{A}}.$$

For each $x \in F$, $\bar{A}x\bar{A}$ is locally closed in the Zariski topology, since it is an orbit of $\bar{A}$ acting on $G(\mathbb{R})/\bar{A}$ [Zim, Corollary 3.1.5]. This means that $\bar{A}x\bar{A} = U \cap E$ where U is Zariski-open and E is Zariski closed. Therefore

$$\overline{Ax\bar{A}} = \overline{U \cap E} \subseteq E \subseteq (U \cap E) \cup (G(\mathbb{R}) \setminus U) = (\bar{A}x\bar{A}) \cup (G(\mathbb{R}) \setminus U).$$

Since $U \subseteq G(\mathbb{R})$ is Zariski open (and $G(\mathbb{R})$ is Zariski connected), $G(\mathbb{R}) \setminus U$ has measure zero, relative to Haar measure μ on $G(\mathbb{R})$ [Mar, Chapter I, Proposition 2.5.3]. Therefore,

$$(2) \quad \mu(\overline{Ax\bar{A}}) = \mu(\bar{A}x\bar{A}).$$

Now we show that $\mu(\bar{A}x\bar{A}) = 0$. Each element of A is semisimple, since each element of Γ is [Mos, Section 11]. Therefore each element of the Zariski closure $\bar{A}$ is also semisimple; in other words, $\bar{A}$ is a torus subgroup. The dimension of $\bar{A}$ as a Lie group is no larger than the absolute rank d' of G (the rank over $\mathbb{C}$ of the Lie algebra of G). The number of positive roots of the complexified Lie algebra of G is at least d' , and the total number of roots is at least $2d'$. Thus the total dimension of the root spaces is at least $2d'$. This means that d' is at most one third of the dimension of G . Thus the dimension of $\bar{A} \times \bar{A}$ is at most two thirds the dimension of G . The map $(a_1, a_2) \mapsto a_1 x a_2$ from $\bar{A} \times \bar{A}$ to $G(\mathbb{R})$ is C^∞ (in fact polynomial). Therefore, by the above dimension count, its image has measure zero. It follows from (2) that $\mu(\bar{A}x\bar{A}) = 0$. However, this contradicts (1), thereby proving the result. $\square$

An immediate consequence of Theorem 1.1 and Theorem 3.2 is

Corollary 3.3. *Let G be a connected semisimple real algebraic group such that $G(\mathbb{R})$ has no compact factors. Let Γ be a uniform lattice subgroup of $G(\mathbb{R})$. Then there exists an abelian subgroup $A < \Gamma$ such that $\text{VN}(A)$ is a masa of $\text{VN}(\Gamma)$ with Pukánszky invariant $\{\infty\}$.*

Proof. This follows immediately from Proposition 3.1. $\square$

Remark 3.4. A similar result was obtained by geometrical methods in [Rob, Theorem 4.6], if Γ is the fundamental group of a compact locally symmetric space M of constant negative curvature and A is generated by the homotopy class of a simple closed geodesic in M .

REFERENCES

- [Bor] A. Borel, *Linear algebraic groups, 2nd edition*, Springer-Verlag, New York, 1991.
- [Dyk] K. Dykema, Two applications of free entropy, *Math. Ann.* **308** (1997), 547–558.
- [DSS] K. J. Dykema, A. M. Sinclair and R. R. Smith, Values of the Pukánszky invariant in free group factors and the hyperfinite factor, *J. Funct. Anal.* **240** (2006), 373–398.
- [GHJ] F. Goodman, P. de la Harpe and V. Jones, *Coxeter graphs and towers of algebras*, Springer-Verlag, New York, 1989.
- [Jol] P. Jolissaint, Operator algebras related to Thompson’s group F , *J. Aust. Math. Soc.* **79** (2005), 231–241.
- [Mar] G. A. Margulis, *Discrete subgroups of semisimple Lie groups*, Springer-Verlag, Berlin, 1991.
- [Mos] G. D. Mostow, *Strong Rigidity of Locally Symmetric Spaces*, Princeton, New Jersey, 1973.
- [NeS] C. Neshveyev, E. Størmer, Ergodic theory and maximal abelian subalgebras of the hyperfinite factor, *J. Funct. Anal.* **195** (2002), 239–261.
- [Po1] S. Popa, Singular maximal abelian $*$ -subalgebras in continuous von Neumann algebras, *J. Funct. Anal.* **50** (1983), 151–166.
- [Po2] S. Popa, Notes on Cartan subalgebras in Type II_1 factors, *Math. Scand.* **57** (1985), 171–188.
- [PrR] G. Prasad and A. S. Rapinchuk, Existence of irreducible $\mathbb{R}$ -regular elements in Zariski-dense subgroups, *Math. Res. Lett.* **10** (2003), 21–32.
- [Rob] G. Robertson, Abelian subalgebras of von Neumann algebras from flat tori in locally symmetric spaces, *J. Funct. Anal.* **230** (2006), 419–431.
- [RoS] G. Robertson and T. Steger, Maximal abelian subalgebras of the group factor of an $\tilde{A}_2$ group, *J. Operator Theory* **36** (1996), 317–334.
- [RSS] G. Robertson, A. M. Sinclair and R. R. Smith, Strong singularity for subalgebras of finite factors, *Int. J. Math.* **14** (2003), 235–258.
- [SS1] A. M. Sinclair and R. R. Smith, The Pukánszky invariant for masas in group von Neumann factors, *Illinois J. Math.* **49** (2003), 325–343.
- [SS2] A. M. Sinclair and R. R. Smith, *Finite von Neumann algebras and masas*, Cambridge University Press, Cambridge, 2008.
- [Whi] S. White, Values of the Pukánszky invariant in McDuff factors, *J. Funct. Anal.* **254** (2008), 612–631.
- [Zim] R. L. Zimmer, *Ergodic Theory and Semisimple Groups*, Birkhäuser, Boston 1985.

SCHOOL OF MATHEMATICS AND STATISTICS, NEWCASTLE UNIVERSITY, NEWCASTLE UPON TYNE NE1 7RU, ENGLAND, U.K.

E-mail address: a.g.robertson@newcastle.ac.uk

STRUTTURA DI MATEMATICA E FISICA, UNIVERSITÀ DI SASSARI, VIA VIENNA 2, 07100 SASSARI, ITALIA

E-mail address: steger@ssmain.uniss.it